

Functions of several variables

Guoning Wu

March 17, 2019

1 The Space \mathbb{R}^m and the Most Important Classes of its Subsets

1.1 The Set \mathbb{R}^m and the distance in it

We make the convention that \mathbb{R}^m denotes the set of ordered m -tuples (x^1, x^2, \dots, x^m) of real numbers $x^i \in \mathbb{R}$.

The function:

$$d(x_1, x_2) = \sqrt{\sum_{i=1}^m (x_1^i - x_2^i)^2}$$

obviously has the following properties:

1. $d(x_1, x_2) \geq 0$;
2. $d(x_1, x_2) = 0 \iff x_1 = x_2$
3. $d(x_1, x_2) = d(x_2, x_1)$;
4. $d(x_1, x_2) \leq d(x_1, x_3) + d(x_3, x_2)$;

A function defined on pairs of points (x_1, x_2) of a set X and possessing the properties 1,2,3,4 is called a **metric or distance on X** .

1.2 Open and Closed Sets in \mathbb{R}^m

Definition 1.1. For each $\delta > 0$, the set

$$B(a, \delta) = \{x \in \mathbb{R}^m \mid d(a, x) < \delta\}$$

is called the ball with center $a \in \mathbb{R}^m$ of radius δ or the δ -neighborhood of the point $a \in \mathbb{R}^m$.

Definition 1.2. A set $G \subset \mathbb{R}^m$ is open in \mathbb{R}^m if for every point $x \in G$ there is a ball $B(a, \delta)$ such that $B(a, \delta) \subset G$.

Definition 1.3. An open set in \mathbb{R}^m containing a given point is called a neighborhood of that point in \mathbb{R}^m .

Definition 1.4. In relation to a set $E \subset \mathbb{R}^m$ a point is

an interior point if some neighborhood of it is contained in E ;

an exterior point if it is a interior point of the complement of E in \mathbb{R}^m ;

a boundary point if it is neither an interior nor an exterior point of E .

2 Limits and Continuity of Functions of Several Variables

2.1 The Limit of a Function

In the next few sections we shall be consider functions $f : X \rightarrow \mathbb{R}^n$ defined on subsets of \mathbb{R}^m .

Definition 2.1. A point $A \in \mathbb{R}^n$ is the **limit of the mapping** $f : X \rightarrow \mathbb{R}^n$ over a base \mathcal{B} in X if for every neighborhood $V(A)$ of the point there exists an element $B \in \mathcal{B}$ of the base whose image $f(B)$ is contained in $V(A)$.

In brief,

$$\lim_{\mathcal{B}} f(x) = A := \forall V(A), \exists B \in \mathcal{B}, f(B) \subset V(A)$$

Examples 1. $\lim_{(x,y) \rightarrow (2,1)} x^2 + xy + y^2 = 7$

Examples 2. $\lim_{(x,y) \rightarrow (0,0)} f(x, y) = 0$, where

$$f(x, y) = \begin{cases} xy \frac{x^2 - y^2}{x^2 + y^2}, & (x, y) \neq (0, 0), \\ 0, & (x, y) = (0, 0) \end{cases}$$

Theorem 2.1. $\lim_{\substack{P \rightarrow P_0 \\ P \in D}} f(P) = A$ 的充要條件是：對於 D 的任意子集 E , 只要 P_0 是 E 的聚點, 就有

$$\lim_{\substack{P \rightarrow P_0 \\ P \in E}} f(P) = A$$

Corollary 2.2. 設 $E_1 \subset D, P_0$ 是 E_1 的聚點，若 $\lim_{\substack{P \rightarrow P_0 \\ P \in E_1}} f(P)$ 不存在，則 $\lim_{\substack{P \rightarrow P_0 \\ P \in D}} f(P)$ 不存在。

Corollary 2.3. 設 $E_1, E_2 \subset D, P_0$ 是它們的聚點，若 $\lim_{\substack{P \rightarrow P_0 \\ P \in E_1}} f(P) = A_1, \lim_{\substack{P \rightarrow P_0 \\ P \in E_2}} f(P) = A_2$ ，但 $A_1 \neq A_2$ ，則 $\lim_{\substack{P \rightarrow P_0 \\ P \in D}} f(P)$ 不存在。

Examples 3. $\lim_{(x,y) \rightarrow (0,0)} f(x,y)$, where

$$f(x,y) = \begin{cases} \frac{xy}{x^2+y^2}, & (x,y) \neq (0,0), \\ 0, & (x,y) = (0,0) \end{cases}$$

2.2 Continuity of a Function of Several Variables and Properties of Continuous Function

Let E be a subset of \mathbb{R}^m and $f : E \rightarrow \mathbb{R}^n$ be a function defined on E with values in \mathbb{R}^n .

Definition 2.2. The function $f : E \rightarrow \mathbb{R}^n$ is a continuous at $a \in E$ if for every neighborhood $V(f(a))$ of the value $f(a)$ that the function assumes at a , there exists a neighborhood $U_E(a)$ of a in E whose image $f(U_E(a))$ is contained in $V(f(a))$.

Thus $f : E \rightarrow \mathbb{R}^n$ is continuous at $a \in E \iff \forall V(f(a)), \exists U_E(a), s.t. f(U_E(a)) \subset V(f(a))$.

It follows from the definition above that the mapping $f : E \rightarrow \mathbb{R}^n$ defined by the relation

$$(x^1, x^2, \dots, x^m) = x \mapsto y = (y^1, y^2, \dots, y^n) = (f^1(x^1, x^2, \dots, x^m), \dots, f^n(x^1, x^2, \dots, x^m))$$

is continuous at a point if and only if each of the function $y^i = f^i(x^1, x^2, \dots, x^m)$ is continuous at that point.

Local properties of continuous functions

a) A mapping $f : E \mapsto \mathbb{R}^n$ is continuous at a point $a \in E$ if and only is $\omega(f; a) = 0$.

b) A mapping $f : E \mapsto \mathbb{R}^n$ is continuous at a point $a \in E$ is bounded in some neighborhood $U_E(a)$ of that point.

c) If the mapping $g : Y \mapsto \mathbb{R}^k$ of the set $Y \subset \mathbb{R}^n$ is continuous at a point $y_0 \in Y$ and the mapping $f : X \mapsto Y$ of the set $X \subset \mathbb{R}^m$ is continuous at a point $x_0 \in X$ and $f(x_0) = y_0$, then the mapping $g \circ f : X \mapsto \mathbb{R}^k$ is defined, and is continuous at $x_0 \in X$.

Global Properties of Continuous function.

a) If a mapping $f : K \mapsto \mathbb{R}^n$ is continuous on a compact set $K \subset \mathbb{R}^m$, then it is uniformly continuous on K .

b) If a mapping $f : K \mapsto \mathbb{R}^n$ is continuous on a compact set $K \subset \mathbb{R}^m$, then it is bounded on K .

c) If a mapping $f : K \mapsto \mathbb{R}^n$ is continuous on a compact set $K \subset \mathbb{R}^m$, then it assumes its maximal and minimal values at some point of K .

d) If a mapping $f : K \mapsto \mathbb{R}^n$ is continuous on a connected set $E \subset \mathbb{R}^m$ and assumes the values $f(a) = A$ and $f(b) = B$ at points $a, b \in E$, then for any C between A and B , there is a point $c \in E$ at which $f(c) = C$.

3 作業

1. 求下列函數的極限

$$(a) \lim_{(x,y) \rightarrow (0,0)} \frac{x^2 y^2}{x^2 + y^2}$$

$$(b) \lim_{(x,y) \rightarrow (0,0)} \frac{1 + x^2 + y^2}{x^2 + y^2}$$

$$(c) \lim_{(x,y) \rightarrow (0,0)} \frac{x^2 + y^2}{\sqrt{1 + x^2 + y^2} - 1}$$

$$(d) \lim_{(x,y) \rightarrow (0,0)} (x + y) \sin \frac{1}{x^2 + y^2}$$

$$(e) \lim_{(x,y) \rightarrow (0,0)} \frac{\sin(x^2 + y^2)}{x^2 + y^2}$$

2. 討論下列函數在點(0, 0)的重極限與累次極限

$$(a) f(x, y) = \frac{y^2}{x^2 + y^2}$$

$$(b) f(x, y) = (x + y) \sin \frac{1}{x} \sin \frac{1}{y}$$

$$(c) f(x, y) = \frac{x^2 y^2}{x^2 y^2 + (x - y)^2}$$

$$(d) f(x, y) = y \sin \frac{1}{x} + x \sin \frac{1}{y}$$

3. 討論下列函數的連續性

$$(a) f(x, y) = \tan(x^2 + y^2)$$

$$(b) f(x, y) = \lfloor x + y \rfloor$$

$$(c) f(x, y) = \begin{cases} \frac{\sin xy}{\sqrt{x^2 + y^2}}, & x^2 + y^2 \neq 0 \\ 0, & x^2 + y^2 = 0 \end{cases}$$

$$(d) f(x, y) = \begin{cases} 0, & x \text{ is an irrational number} \\ y, & x \text{ is a rational number} \end{cases}$$